

Information sheet for customer

Crude drug Extract Preparations

2nd class OTC drugs

JINSENSAN

- ◇JINSENSAN is a medicine consisting of 15 kinds of crude drugs.
- ◇Most cases of cystitis are inflammation caused by bacteria. "JINSENSAN" contains an antibacterial crude drug 'Bearberry Leaf' that works against the causative bacteria.

PRECAUTIONS

Who should seek consultation

- The following persons should consult a physician, pharmacist, or registered distributor before taking this medicine.
 - Patients undergoing medical treatment from a physician
 - Pregnant women and women suspected of being pregnant
 - Persons who have gastrointestinal weakness
 - Patients who have had allergic symptoms to drugs, etc
 - Persons with the following symptoms: loss of appetite, nausea·vomiting
- If the following symptoms are observed after taking this medicine, these may be adverse reactions, so immediately discontinue the use of this medicine, and show this leaflet to your physician, pharmacist, or registered distributor for a consultation.

Affected areas	Symptoms
Skin	Rash·redness, itching
Gastrointestinal system	Loss of appetite, stomach discomfort, nausea·vomiting

The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid.

Symptom's name	Symptoms
Mesenteric phlebosclerosis	Abdominal pain, diarrhea, constipation, abdominal distension, etc. may occur repeatedly with long-term oral administration.

- The following symptoms may be observed after taking this medicine. If these symptoms persist or worsen, discontinue the use of this medicine, and show this leaflet to your physician, pharmacist, or registered distributor for a consultation.

Diarrhea

- When symptoms do not improve even after taking the medicine for about 1 month, stop taking this medicine and consult a physician, pharmacist or registered distributor, being sure to take this leaflet with you.
- If you take this medicine for a long period, consult a physician, pharmacist, or registered distributor.

[INDICATION]

Nephritis, Nephrosis, Pyelonephritis, Cystitis, Edema, Decreased excretion of urine

[DOSAGE•ADMINISTRATION]

Take the following dosages with water or warm water between meals. (on an empty stomach)

Age	One dosage	Daily dosage
Adult	1 packet	3 times
From 8 to 15years	1/2 packet	
From 4 to 7 years	1/3 packet	
Less than 4 years	Do not use	

※When allowing children to take this medicine, a guardian must be present to watch and tell such them how to take it.

[INGREDIENT•COMPOSITION]

20 packets contain extract of the following mixed crude drugs. (1.5g per 1 packet)

Alisma Rhizome	6.0 g	Cinnamon Bark	6.0 g	Rehmannia Root	6.0 g
Bearberry Leaf	10.0g	Chuling	6.0 g	Artemisia Capillaris Flower	5.0 g
Sinomenium Stem	6.0 g	Plantago Seed	6.0 g	Hoelen	8.0 g
Gardenia Fruit	5.0 g	Sambucus Rasemosa Wood	8.0 g	Imperata Rhizome	6.0 g
Atractylodes Lancea Rhizome	7.0 g	Peony Root	7.0 g	Catalpa Fruit	8.0 g

Inactive ingredients: Carmellose Ca, Silicic anhydride

[PRECAUTIONS FOR STORAGE AND HANDLING]

- (1) Store in a cool, low humidity place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer this medicine to a different container.
(It may lead to misuse or alter the quality of the drug.)
- (4) Take this medicine within 2days, when you take separately.
- (5) Do not take this medicine after the expiration date.

Mayado pharmaceutical company
<https://www.mayado.jp/about/>