

Information sheet for customer

Crude drug Extract Preparations

2nd class OTC drugs

TŌKAIJYO

- ☆ TŌKAIJYO is a medicine consisting of 10 kinds of crude drugs. It has the function of gently lowering blood glucose.
- ☆ Improves various symptoms caused by diabetes (dry mouth, frequent urination, polyuria).
- ☆ This medicine is sugar-coated tablet easy to take.

Precautions

Who should seek consultation

1. The following persons should consult a physician, pharmacist, or registered distributor before taking this medicine.

- (1) Persons undergoing medical treatment from a physician
- (2) Pregnant women and women suspected of being pregnant
- (3) Persons who have gastrointestinal weakness.
- (4) Elderly people
- (5) Persons with the following symptoms:
edema, loss of appetite, nausea · vomiting
- (6) Persons diagnosed as having the following:
hypertension, heart disease, kidney disease

2. The following symptoms may be adverse reactions to this medication.

If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist, or registered distributor. Take the instruction leaflet with you.

Related areas	Symptoms
Skin	rash · redness, itching
Gastrointestinal system	loss of appetite, gastric distress, nausea · vomiting
Others	excitement, sleep loss, hypertension

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

Symptom's name	Symptoms
Pseudoaldosteronism, myopathy	In addition to listlessness, numbness, stretched feeling, and stiffness of hands and feet, feeling of weakness and muscular pain will appear and gradually progress.

3. The following symptom may occur after administration of this medicine. If the symptoms persist or increase, stop taking this medicine immediately and consult a physician, pharmacist, or registered distributor. Take the instruction leaflet with you.

Diarrhoea

- 4. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 month. Take the leaflet with you.
- 5. If you take this medicine regularly for a long period, consult a physician, pharmacist or registered distributor.

indications

Diabetes

Dosage and administration

Take the following dosages with water or warm water before or between meals.

(on an empty stomach)

Age	One dosage	Daily dosage
15 years and over	3 to 5 tablets	3 to 5 times a day
Less than 15 years	Do not use	

Ingredient and Composition

120 tablets contain the following ingredients.

Ingredient	Composition	Ingredient	Composition
Ophiopogon Root	12.0 g	Trichosanthes Root	12.0 g
Ginseng	8.0 g	Poria Sclerotium	12.0 g
Pueraria Root	12.0 g	Schisandra Fruit	6.0 g
Rehmannia Root	12.0 g	Glycyrrhiza	6.0 g
Anemarrhena Rhizome	10.0 g	Aralia Elata Root Bark	10.0 g

Inactive ingredients: Carboxymethyl cellulose, Aluminium Hydroxide, Hydroxypropylcellulose, Magnesium aluminometasilicate, Cellulose, Magnesium stearate, Sodium copper chlorophyllin, Gum Arabic, Gelatin, White soft sugar, Calcium carbonate, Talc, Shellac, Hypromellose, Macrogol, Carnauba wax

Precautions for storage and handling

- ① Store in a cool, dry place free from direct sunlight.
- ② Keep all medication out of reach of children.
- ③ Do not transfer the medicine to a different container.
(It may lead to misuse or alter the quality of the drug.)
- ④ Close the bottle cap tightly after use to prevent deterioration of tablet quality from moisture.
Do not touch this medicine with wet hands.
- ⑤ Discard a packing in a bottle after opening because it is used to prevent the damage of tablets during the transportation.
- ⑥ Fill in the “Opening date” of the box and bottle after opening.
- ⑦ Take this medicine as soon as possible to keep quality.
- ⑧ Do not take this medicine after the expiration date.